

Technical Bulletin

Genes that fit *your* farm.

SeCan

Canada's Seed Partner

AAC Tisdale Canada Western Red Spring Wheat



Progress Through Research
Le progrès grâce à la recherche

Description:

AAC Tisdale is an awned CWRS variety with consistently low DON accumulation, MR rating to FHB and high grain protein. AAC Tisdale has good lodging tolerance, improved grain yield and good overall disease package. AAC Tisdale is well suited to all growing regions of western Canada.

Parentage: Somerset/BW865/Waskada

Strengths:

- MR to FHB and low DON accumulation
- Grain yield 104% of the mean of the checks, over all sites in 2013 to 2015 Coop Registration trials
- High protein potential based on provincial regional variety trials
- Improved lodging compared to the checks
- Resistant to stem and leaf rust
- MR to common bunt and loose smut
- Large seed size

Neutral Traits:

- Later maturing than the checks (1 day earlier than AC® Carberry)
- 7cm taller than AC® Carberry

Weaknesses:

- Moderately susceptible to stripe rust

Breeder:

Dr. Richard Cuthbert
Semiarid Prairie Agricultural Research Centre
Agriculture and Agri-Food Canada
Swift Current, SK

PBR 91 Pending

2013-2015 Parkland Wheat Cooperative Trials - Registration Data

Variety	Yield (% of Check Mean)	Maturity (days)	Lodging 1 = erect 9 = flat	Height (cm)	Test Weight (kg/kl)	Kernel Weight (mg/kernel)	Grain Protein (%)	Falling Number
AC Splendor	94.5	95.9	2.7	98	78.0	34.9	14.5	405
PT472	104.7	96.8	2.2	98	80.8	36.3	14.2	445
Parata	100.8	97.7	2.2	93	80.4	34.8	14.5	400
AAC Tisdale	103.6	100.1	1.9	93	80.5	37.5	14.5	395

'AC' is an official mark used under license from Agriculture & Agri-Food Canada

2019 Seed Manitoba - Wheat Comparison

Variety	Site Years Tested	Yield bu/ac	Protein %	Maturity +/- 99 days	Height +/- 81cm	Spike Awned	Resistance to:								
							Lodging	Sprouting	Loose Smut	Bunt	Leaf Spot	Stem Rust	Leaf Rust	Stripe Rust	FHB
AAC Brandon	60	69	14.2	+2	0	Y	VG	P	MR	S	I	R	R	MR	MR
AC® Carberry	110	66	14.5	+2	0	Y	VG	F	MR	R	MS	MR	R	MR	MR
AC® Cardale	28	67	14.5	+2	+3	Y	VG	G	I	MR	MS	R	R	S	MR
AAC Elie	44	68	14.3	+2	-3	Y	VG	F	I	I	I	R	R	MR	I
AAC Redwater	43	64	14.5	-2	+8	Y	G	VG	MS	I	MS	R	R	MR	I
CDC Bradwell	33	66	14.4	+3	+5	Y	VG	F	MR	R	MS	MR	R	MS	I
Parata	6	69	14.2	-2	+13	Y	G	---	MR	S	I	R	MR	MR	I
AAC Tisdale	32	67	15.1	+1	+8	Y	G	F	MR	MR	MS	R	R	S	MR

Lodging Ratings: F=Fair; G=Good; VG=Very Good

Disease Ratings: R=Resistant; MR=Moderately Resistant; I=Intermediate; MS=Moderately Susceptible; S=Susceptible

2019 Varieties of Grain Crops for Saskatchewan – Wheat Comparison

Variety	Years Tested	Yield as % of Carberry			Protein	Resistance to:									Relative Maturity (days)	Head Awedness	Seed Weight (mg)	Test Weight (kg/hl)	Height (cm)
		Area 1 & 2	Area 3 & 4	Irrigation		Lodging	Sprouting	Stem Rust	Leaf Rust	Stripe Rust	Loose Smut	Bunt	Leaf Spot	FHB					
AC® Carberry	6	100	100	100	14.6	VG	F	MR	R	MR	MR	R	MS	MR	99	Y	35.7	80.3	83
AAC Brandon	5	106	106	---	-0.4	G	P	R	R	MR	MR	S	I	MR	0	Y	+0.1	-0.1	0
AAC Elie	5	105	105	---	-0.2	G	F	R	R	MR	I	I	I	I	0	Y	-0.1	0.0	-2
AAC Redwater	5	102	101	---	+0.1	F	VG	R	R	MR	MS	I	MS	I	-3	Y	-3.5	-1.3	+8
AC® Stettler	6	105	107	100	+0.2	F	G	MR	MS	MR	R	MR	MS	MS	-1	Y	-0.6	-0.4	+8
CDC Bradwell	5	101	108	---	0.0	VG	F	MR	R	MS	MR	R	MS	I	0	Y	-2.0	+0.6	+8
Parata	2	98	106	---	+0.3	F	F	R	MR	MR	MR	S	I	I	-2	Y	-2.0	-0.1	+11
AAC Tisdale	3	100	109	---	+0.7	F	F	R	R	S	MR	MR	MS	MR	-2	Y	+0.8	-0.3	+8

G=Good; VG=Very Good; F=Fair; P=Poor; VP=Very Poor

2019 Alberta Seed Guide - CWRs Wheat Comparison

Variety	Overall Yield		Test Yield Category			Maturity Rating	Protein %	Test Weight (lb/bu)	Kernel Weight g/1000	Height (cm)	Resistance to:		Disease Resistance:						
	All Sites	Station years of testing	Low < 55 bu/ac	Med 55 - 80 bu/ac	High >80 bu/ac						Ldg.	Sprout	Loose Smut	Bunt	Stripe Rust	Leaf Spot	FHB		
			Yield as % of AC® Carberry																
AC® Carberry bu/ac	67		45	65	92														
AC® Carberry	100		100	100	100	L	14.0	63	39	79	VG	F	MR	R	MR	MS	MR		
AAC Brandon	106	21	XX	103	110	M	-0.2	63	40	80	VG	P	MR	S	MR	I	MR		
AAC Redwater	96	41	91	101	98	E	0.0	64	35	87	G	VG	MS	I	MR	MS	I		
AC® Stettler	106	82	105	105	104	M	0.0	63	39	85	G	G	R	MR	I	MS	MS		
CDC Bradwell	101	42	97	105	101	L	-0.3	63	38	84	VG	F	MR	R	MS	MS	I		
Parata	97	31	XX	98	96	E	+0.7	64	39	87	F	F	MS	S	MR	I	I		
AAC Tisdale	101	45	100	101	102	M	+0.4	63	42	88	F	F	MR	MR	S	MS	MR		

VG = Very Good; G = Good; F = Fair; P = Poor; VP = Very Poor; Disease Ratings: R=Resistant; MR=Moderately Resistant; I=Intermediate; MS=Moderately Susceptible; S=Susceptible

For more information, call 1-800-665-7333 or visit www.secan.com